

## FINITE-TIME STABILIZATION OF A NETWORK OF STRINGS

FATIHA ALABAU-BOUSSOIRA

Institut Elie Cartan de Lorraine, UMR 7502  
Université de Lorraine  
57045 Metz, France

VINCENT PERROLLAZ\* AND LIONEL ROSIER\*\*

\*Laboratoire de Mathématiques et Physique Théorique  
Université de Tours, UFR Sciences et Techniques  
Parc de Grandmont  
37200 Tours, France

\*\*Centre Automatique et Systèmes  
MINES ParisTech, PSL Research University  
60 Boulevard Saint-Michel  
75272 Paris Cedex 06, France

(Communicated by Enrique Zuazua)

**ABSTRACT.** We investigate the finite-time stabilization of a tree-shaped network of strings. Transparent boundary conditions are applied at all the external nodes. At any internal node, in addition to the usual continuity conditions, a modified Kirchhoff law incorporating a damping term  $\alpha u_t$  with a coefficient  $\alpha$  that may depend on the node is considered. We show that for a convenient choice of the sequence of coefficients  $\alpha$ , any solution of the wave equation on the network becomes constant after a finite time. The condition on the coefficients proves to be sharp at least for a star-shaped tree. Similar results are derived when we replace the transparent boundary condition by the Dirichlet (resp. Neumann) boundary condition at one external node. Our results lead to the finite-time stabilization even though the systems may not be dissipative.

**1. Introduction.** Solutions of certain ODE  $\dot{x} = f(x)$  may reach the equilibrium state in finite time. This phenomenon, when combined with the stability, was termed *finite-time stability* in [4, 11].

A *finite-time stabilizer* is a feedback control for which the closed-loop system is finite-time stable around some equilibrium state. In some sense, it satisfies an exact controllability objective with a control in feedback form. On the other hand, a finite-time stabilizer may be seen as an exponential stabilizer yielding an arbitrarily large decay rate for the solutions to the closed-loop system. Indeed, any solution of the closed-loop system can be estimated as

$$\|x(t)\| \leq h(\|x_0\|)\mathbf{1}_{[0,T]}(t) \leq h(\|x_0\|)e^{-\lambda(t-T)}$$

---

2010 *Mathematics Subject Classification.* Primary: 93D15; Secondary: 34B45, 35L05.

*Key words and phrases.* Finite-time stabilization, network, wave equation, transparent boundary condition, Kirchhoff law, Riemann invariant.

where  $h(\delta) \rightarrow 0$  as  $\delta \rightarrow 0$ , and  $\lambda > 0$  is arbitrarily large. This explains why some efforts were made in the last decade to construct finite-time stabilizers for controllable systems, including the linear ones. See [15] for some recent developments and up-to-date references, and [3] for some connections with Lyapunov theory.

To the best knowledge of the authors, the analysis of the finite-time stabilization of PDE is not developed yet. However, since [14], it is well-known that solutions of the wave equation on certain bounded domains may disappear when using *transparent* boundary conditions. For instance, the solution of the 1-D wave equation

$$u_{tt} - c^2 u_{xx} = 0, \quad \text{in } (0, L) \times (0, T), \quad (1)$$

$$cu_x(L, t) = -u_t(L, t), \quad \text{in } (0, T), \quad (2)$$

$$cu_x(0, t) = u_t(0, t), \quad \text{in } (0, T), \quad (3)$$

$$(u(0), u_t(0)) = (u^0, u^1), \quad \text{in } (0, L), \quad (4)$$

is finite-time stable in the space  $\{(u, v) \in H^1(0, L) \times L^2(0, L); c(u(0) + u(L)) + \int_0^L v(x)dx = 0\}$ , with  $T = L/c$  as extinction time (see e.g. [12, Theorem 0.5] for the details.) The condition (2) is “transparent” in the sense that a wave  $u(x, t) = f(x - ct)$  traveling to the right satisfies (2) and leaves the domain at  $x = L$  without generating any reflected wave. Note that the solution issued from any state  $(u^0, u^1) \in H^1(0, L) \times L^2(0, L)$  is not necessarily vanishing, but constant, for  $t \geq L/c$ . Note also that if we replace (3) by the boundary condition  $u(0, t) = 0$  (or  $u_x(0, t) = 0$ ), then a finite-time extinction still occurs (despite the fact that waves bounce at  $x = 0$ ) with an extinction time  $T = 2L/c$ . We refer to [5] for the analysis of the finite-time extinction property for a nonhomogeneous string with a viscous damping at one extremity, to [8] for the finite-time stabilization of a string with a moving boundary, to [16] (resp. [17]) for the finite-time stabilization of a system of conservation laws on an interval (resp. on a tree-shaped network).

The finite-time stability of (1)-(4) is easily established when writing (1) as a system of two transport equations

$$d_t + cd_x = 0,$$

$$s_t - cs_x = 0.$$

where  $d := u_t - cu_x$  and  $s := u_t + cu_x$  stand for the Riemann invariants for the wave equation written as a first order hyperbolic system. The boundary conditions (2) and (3) yield  $d(0, t) = s(L, t) = 0$  (and hence  $d(\cdot, t) = s(\cdot, t) = 0$  for  $t \geq L/c$ ), while the boundary conditions (2) and  $u(0, t) = 0$  yield  $s(L, t) = 0$  and  $d(0, t) = -s(0, t)$  (and hence  $s(\cdot, t) = 0$  for  $t \geq L/c$  and  $d(\cdot, t) = 0$  for  $t \geq 2L/c$ ).

The stabilization of networks of strings has been considered in e.g. [1, 2, 7, 9, 10, 19, 22]. In [10], the authors considered a star of vibrating strings, and derived the finite-time stability (resp. the exponential stability) when transparent boundary conditions are applied at all external nodes (resp. at all external nodes but one, which is changing as times proceeds). For a more general network, we guess that the finite time stability cannot hold without the introduction of additional feedback controls at the internal nodes. Indeed, it is proved here that for a bone-shaped tree, if the feedback controls are applied only at the external nodes, then the finite-time stability fails.

The aim of this paper is to investigate the *finite-time* stabilization of a *tree-shaped* network of strings. At each internal node  $n$  connecting  $k$  edges, we assume that the usual continuity condition hold:

$$u_i(n, t) = u_j(n, t), \quad \text{for all indices } i, j \text{ of edges having } n \text{ as one end,} \quad (5)$$

while the usual Kirchhoff law is modified by incorporating a *damping term* inside:

$$\sum_i c_i u_{i,x}(n, t) - c_{i_0} u_{i_0,x}(n, t) = -\alpha(n) u_t(n, t). \quad (6)$$

In (6), the sum is over the indices  $i$  of the edges having  $n$  as initial point,  $i_0$  denotes the index of the edge having  $n$  as final point (the tree being oriented),  $\alpha(n) \in \mathbb{R}$  is a coefficient depending on the node  $n$ , and  $u(n, t) := u_i(n, t)$  (for any  $i$ ). The case  $\alpha = 0$  corresponds to the usual (conservative) Kirchhoff law.

Note that we can assume without loss of generality that the length of each edge is one, by scaling the variable  $x$  and the coefficient  $c_i$  along each edge.

Even if the finite-time stabilization of  $2 \times 2$  hyperbolic systems on tree-shaped networks was already considered in [17] (and applied to the regulation of water flows in networks of canals, with  $k - 1$  controls at any node connecting  $k$  canals), the novelty (and difficulty) here comes from the fact that only *one* control is applied at each internal node. The present work can be seen as a first step in the understanding of the finite-time stabilization of systems of conservation laws with few controls. In practice, on account of possible budget limitations, it is important that the number of interior controls be chosen as small as possible.

A natural guess is that the finite-time stability cannot hold if one can find in the tree a pair of adjacent nodes that are free of any control, because of the (partial but standing) bounces of waves at these nodes. This conjecture will be demonstrated here for a star-shaped tree and a bone-shaped tree.

Actually, we shall prove that the finite-time stabilization can be achieved for a very particular choice of the coefficient  $\alpha$  at each internal node. One of the main results proved in this paper is the following

**Theorem 1.1.** *Consider any tree-shaped network of strings, with transparent boundary conditions at all the external nodes but one (the root), where a homogeneous Dirichlet boundary condition is applied, and with the continuity conditions and the modified Kirchhoff law at the internal nodes. If at each internal node  $n$  connecting  $k$  edges we have*

$$\alpha(n) = k - 2, \quad (7)$$

*then each solution of the wave equation on the network becomes null after some finite time. Conversely, the condition (7) is also necessary for the finite-time stability of the system on a star-shaped tree.*

Similar results will be obtained when replacing at the root the homogeneous Dirichlet boundary condition by the homogeneous Neumann boundary condition or by the transparent boundary condition. The necessity of the condition  $\alpha = k - 2$  for the finite-time stability of a star-shaped tree is obtained by doing some explicit computation of the discrete spectrum of the underlying operator. The same approach gives for a bone-shaped tree a necessary and sufficient condition for the finite-time stability, which differs slightly from those stated in Theorem 1.1.

The condition  $\alpha(n) = k - 2$  can be interpreted as follows: the wave emanating from  $n$  along any edge is the sum of the waves arriving at  $n$  along the  $k - 1$  others edges (see below (34)). The node  $n$  is *transparent* in the sense that there is transmission without attenuation nor reflection of all the waves that reach it.

The well-posedness of the system is also completely characterized in terms of the coefficients  $\alpha$  (see below Theorem 2.1). We note that the well-posedness of

the system may hold even if the system is not dissipative, and that our finite-time stability results hold in a class of systems that are not necessarily dissipative.

Finally, we point out that the recent papers [18, 21] contain some results similar to ours. (Note that the finite-time stability is termed *super-stability* in [18, 21].) The paper [21] is concerned with star-shaped trees, while the paper [18] considers very general networks. When comparing the results in [18] with ours, we observe the following:

1. The wave equation in [18] takes the form

$$\rho_i u_{i,tt} - T_i u_{i,xx} = 0, \quad \text{in } (0, 1) \times (0, T) \quad (8)$$

where  $T_i, \rho_i > 0$  are given coefficients, while the modified Kirchhoff law at an internal node of a tree is written

$$\sum_i T_i u_{i,x}(n, t) - T_{i_0} u_{i_0,x}(n, t) = -\alpha(n) u_t(n, t). \quad (9)$$

Note that  $(T_i, \rho_i) := (c_i, c_i^{-1})$  is a possible choice (among others) to yield (12) (see below) on the edge  $i$ .

2. The well-posedness of the complete system is derived in [18, Proposition 2.1] under the *sufficient* condition  $\alpha(n) \leq 0$  for all internal nodes  $n$ , which ensures that the system is *dissipative*, while a *sharp* condition is given in our Theorem 2.1.
3. When  $(T_i, \rho_i) = (c_i, c_i^{-1})$  for all  $i$ , the condition (3.7) in [18, Theorem 3.2] which gives the values of  $\alpha(n)$  at all the internal nodes ensuring the finite-time stability is the same as our condition (92) in Theorem 4.1. However, it is also assumed in [18, Theorem 3.2] that

$$\alpha(n) \leq 0 \quad \text{for all internal nodes } n. \quad (10)$$

This imposes some restrictions on the coefficients  $T_i, \rho_i$  or on the shape of the tree. For instance, for uniform data ( $T_i = \rho_i = 1$  for all  $i$ ), then the condition (10) (with  $\alpha_n$  as in (92)) is satisfied only if  $k_n = 2$  for all internal nodes  $n$ , i.e. only if the tree is a *line*, while our Theorem 4.1 is valid for *any tree*. On the other hand, by choosing other pairs of positive coefficients  $(T_i, \rho_i)$  with  $T_i/\rho_i = c_i^2$ , Theorem 3.2 in [18] gives modified Kirchhoff laws (9) different from (6) that may lead to the finite-time stability.

4. Our approach is based upon the analysis of the Riemann invariants and the propagation of waves, while [18] is based upon spectral theory.

The paper is outlined as follows. In Section 2, we provide a sharp condition on the coefficients  $\alpha(n)$  for the system to be well-posed. It is obtained by expressing the conditions (5)-(6) at the internal nodes in terms of the Riemann invariants. In Section 3, we consider the particular case of a tree with two internal nodes to obtain sharp results for the finite-time stability. In Section 4, we derive the finite-time stability results when the coefficients  $\alpha$  are chosen as in (7). Finally, we prove in Section 5 the necessity of that condition for tree-shaped networks.

**2. Well-posedness.** We introduce some notations inspired by [6]. Let  $\mathcal{T}$  be a tree, whose *vertices* (or *nodes*) are numbered by the index  $n \in \mathcal{N} = \{0, \dots, N\}$ , and whose *edges* are numbered by the index  $i \in \mathcal{I} = \{1, \dots, N\}$ . We choose a simple vertex (i.e. an external node), called the *root* of  $\mathcal{T}$  and denoted by  $\mathcal{R}$ , and which corresponds to the index  $n = 0$ . The edge containing  $\mathcal{R}$  has  $i = 1$  as index, and its other endpoint has for index  $n = 1$ . We choose an orientation of the edges in the tree such that  $\mathcal{R}$  is

the “first” encountered vertex. The *depth*  $d$  of the tree is the number of generations ( $d = 1$  for a tree reduced to a single edge,  $d = 2$  for a star-shaped tree, etc.) Once the orientation of the tree is chosen, each point of the  $i$ -th edge (of length 1) is identified with a real number  $x \in [0, 1]$ . The points  $x = 0$  and  $x = 1$  are termed the *initial point* and the *final point* of the  $i$ -th edge, respectively. Renumbering the edges if needed, we can assume that the edge of index  $i$  has as final point the vertex with the (same) index  $n = i$  for all  $i \in \mathcal{I}$ . (See Figure 1.) The set of indices of




FIGURE 1. A tree with 14 nodes, a depth equal to 5, with simple nodes  $\mathcal{N}_S = \{0, 4, 8, 9, 10, 11, 12, 13\}$  and multiple nodes  $\mathcal{N}_M = \{1, 2, 3, 5, 6, 7\}$ .

simple and multiple nodes are denoted by  $\mathcal{N}_S$  and  $\mathcal{N}_M$ , respectively.

For  $n \in \mathcal{N}_M$  we denote by  $\mathcal{I}_n$  the set of indices of those edges having the vertex of index  $n$  as initial point. As we consider a network of strings whose constants  $c_i$  may vary from one edge to another one, the case  $\#\mathcal{I}_n = 1$  (one child) is possible. The number of edges having the vertex of index  $n$  as one of their extremities, also termed the *degree* of the vertex  $n$ , is

$$k_n := \#\mathcal{I}_n + 1 \geq 2. \quad (11)$$

We consider the following system

$$u_{i,tt} - c_i^2 u_{i,xx} = 0, \quad t > 0, 0 < x < 1, i \in \mathcal{I} \quad (12)$$

$$(u_i(\cdot, 0), u_{i,t}(\cdot, 0)) = (u_i^0, u_i^1), \quad i \in \mathcal{I} \quad (13)$$

with the following boundary conditions

$$c_n u_{n,x}(1, t) = -u_{n,t}(1, t), \quad t > 0, n \in \mathcal{N}_S \setminus \{0\}, \quad (14)$$

$$\sum_{i \in \mathcal{I}_n} c_i u_{i,x}(0, t) - c_n u_{n,x}(1, t) = -\alpha_n u_{n,t}(1, t), \quad t > 0, n \in \mathcal{N}_M, \quad (15)$$

$$u_i(0, t) = u_n(1, t), \quad t > 0, n \in \mathcal{N}_M, i \in \mathcal{I}_n, \quad (16)$$

where the sequence  $(\alpha_n)_{n \in \mathcal{N}_M}$  is still to be defined. For the boundary condition at the root  $\mathcal{R}$ , we shall consider one of the following conditions

$$u_1(0, t) = 0, \quad t > 0 \quad (\text{Dirichlet boundary condition}); \quad (17)$$

$$u_{1,x}(0, t) = 0, \quad t > 0 \quad (\text{Neumann boundary condition}); \quad (18)$$

$$c_1 u_{1,x}(0, t) = u_{1,t}(0, t), \quad t > 0 \quad (\text{Transparent boundary condition}). \quad (19)$$

Let

$$\mathcal{H} = \{(u_i, v_i)_{i \in \mathcal{I}} \in \prod_{i \in \mathcal{I}} [H^1(0, 1) \times L^2(0, 1)]; u_i(0) = u_n(1) \forall n \in \mathcal{N}_M, \forall i \in \mathcal{I}_n\}$$

and  $\mathcal{H}_0 = \{(u_i, v_i)_{i \in \mathcal{I}} \in \mathcal{H}; u_1(0) = 0\}$ . The spaces  $\mathcal{H}$  and  $\mathcal{H}_0$  are Hilbert spaces when endowed with the scalar product

$$((u_i, v_i)_{i \in \mathcal{I}}, (\tilde{u}_i, \tilde{v}_i)_{i \in \mathcal{I}})_{\mathcal{H}} := \sum_{i \in \mathcal{I}} \int_0^1 (u_i(x)\tilde{u}_i(x) + u_{i,x}(x)\tilde{u}_{i,x}(x) + v_i(x)\tilde{v}_i(x)) dx.$$

Replacing  $u_{i,t}$  by  $v_i$  and dropping the variable  $t$ , conditions (14) - (19) may be rewritten respectively as

$$c_n u_{n,x}(1) = -v_n(1), \quad n \in \mathcal{N}_S \setminus \{0\}, \quad (20)$$

$$\sum_{i \in \mathcal{I}_n} c_i u_{i,x}(0) - c_n u_{n,x}(1) = -\alpha_n v_n(1), \quad n \in \mathcal{N}_M, \quad (21)$$

$$u_i(0) = u_n(1), \quad n \in \mathcal{N}_M, i \in \mathcal{I}_n, \quad (22)$$

$$u_1(0) = 0, \quad (23)$$

$$u_{1,x}(0) = 0, \quad (24)$$

$$c_1 u_{1,x}(0) = v_1(0). \quad (25)$$

If  $t \in \mathbb{R}^+ \rightarrow (u_i, v_i)_{i \in \mathcal{I}} \in \mathcal{D}(A_T)$  is continuous (for the definition of  $\mathcal{D}(A_T)$ , see below), using  $v_i = u_{i,t}$ , (22) and (23) we obtain

$$v_i(0) = v_n(1), \quad n \in \mathcal{N}_M, i \in \mathcal{I}_n, \quad (26)$$

$$v_1(0) = 0. \quad (27)$$

Introduce the operator  $A_D$ ,  $A_N$  and  $A_T$  defined as

$$A_D((u_i, v_i)_{i \in \mathcal{I}}) = (v_i, c_i^2 u_{i,xx})_{i \in \mathcal{I}},$$

$$A_N((u_i, v_i)_{i \in \mathcal{I}}) = (v_i, c_i^2 u_{i,xx})_{i \in \mathcal{I}},$$

$$A_T((u_i, v_i)_{i \in \mathcal{I}}) = (v_i, c_i^2 u_{i,xx})_{i \in \mathcal{I}},$$

with respective domains

$$\mathcal{D}(A_D) = \{(u_i, v_i)_{i \in \mathcal{I}} \in \prod_{i \in \mathcal{I}} [H^2(0, 1) \times H^1(0, 1)]; (20) - (22), (23)$$

$$\text{and (26) - (27) hold}\} \subset \mathcal{H}_0,$$

$$\mathcal{D}(A_N) = \{(u_i, v_i)_{i \in \mathcal{I}} \in \prod_{i \in \mathcal{I}} [H^2(0, 1) \times H^1(0, 1)]; (20) - (22), (24)$$

$$\text{and (26) hold}\} \subset \mathcal{H},$$

$$\mathcal{D}(A_T) = \{(u_i, v_i)_{i \in \mathcal{I}} \in \prod_{i \in \mathcal{I}} [H^2(0, 1) \times H^1(0, 1)]; (20) - (22), (25)$$

$$\text{and (26) hold}\} \subset \mathcal{H}.$$

The main result in this section is concerned with the well-posedness of system (12)-(16) and (17) (or (18), or (19)).

**Theorem 2.1.** *Let  $\mathcal{T}$  be a tree and let  $(\alpha_n)_{n \in \mathcal{N}_M}$  be a given family of real numbers. Then  $A_T$  generates a strongly continuous semigroup of operators on  $\mathcal{H}$  if, and only if,*

$$\alpha_n \neq k_n \quad \forall n \in \mathcal{N}_M, \quad (28)$$

where  $k_n$  is the degree of the vertex  $n$  defined in (11). The same conclusion holds for  $A_N$  on  $\mathcal{H}$  (resp. for  $A_D$  on  $\mathcal{H}_0$ ).

**Remark 1.** (1) As a consequence of Theorem 2.1, system (12)-(16) and (19) is *not well-posed* when  $\alpha_n = k_n$  for some  $n \in \mathcal{N}_M$ . The initial data has to fulfill some compatibility conditions for the solution to exist. Consider the simplest case of a tree with two edges ( $N = 2$ ) and  $c_1 = c_2 = 1$ . From (35) (see below), we have

$$v_1^0(1-t) - u_{1,x}^0(1-t) + v_2^0(t) + u_{2,x}^0(t) = 0 \quad \text{for a.e. } t \in (0, 1). \quad (29)$$

This is a *necessary* condition on the initial data for the condition (15)-(16) to be satisfied. In other words, for initial data not satisfying the condition (29), there is *no solution* to (12)-(16) and (19).

(2) Furthermore, even for initial data satisfying (29), the corresponding solutions are *not unique* for positive times. Indeed, taking 0 as initial data, and picking any function  $h \in C^1(\mathbb{R})$  with  $h(t) = 0$  for  $t \leq 0$ , then the trajectory candidate

$$\begin{aligned} u_1(x, t) &= h(t+x-1), & 0 \leq x \leq 1, t \geq 0, \\ u_2(x, t) &= h(t-x), & 0 \leq x \leq 1, t \geq 0 \end{aligned}$$

satisfies (12)-(16) and (19) with  $\alpha_1 = k_1 = 2$ . Thus, there is no uniqueness of trajectories.

(3) On the other hand, system (12)-(16) and (19) is *well-posed* when  $\alpha_n \neq k_n$  for all  $n \in \mathcal{N}_M$ . Note that the system is not necessarily dissipative, i.e. the energy is not necessarily nonincreasing (even if the system is finite-time stable). Consider for instance a star-shaped tree with 3 edges ( $N = 3$ ), and  $c_1 = c_2 = c_3 = 1$  for simplicity. Pick  $\alpha_1 := k_1 - 2 = 1$ . (According to Theorem 4.1, the system is finite-time stable to constant functions.) Then a direct computation gives for any solution  $(u_i, v_i)_{1 \leq i \leq 3}$  of (12)-(16) and (19)

$$\frac{d}{dt} \frac{1}{2} \sum_{i=1}^3 \int_0^1 (u_{i,t}^2 + u_{i,x}^2) dx = u_{1,t}^2(1, t) - (u_{1,t}^2(0, t) + u_{2,t}^2(1, t) + u_{3,t}^2(1, t)).$$

This expression is *positive* for  $t > 0$  small enough if the initial data  $(u_i^0, v_i^0)_{1 \leq i \leq 3} \in \mathcal{D}(A_T)$  is chosen so that

$$|v_1^0(1)|^2 > |v_1^0(0)|^2 + |v_2^0(1)|^2 + |v_3^0(1)|^2.$$

In that case, the energy may increase before reaching the value 0 in finite time.

*Proof.* We sketch the proof only for  $A_T$ . We need a preliminary result about the Riemann invariants around an internal node. Consider any internal node connecting edges whose indices range over  $\{1, \dots, k\}$  (to simplify the notations). Consider any solution of (12) satisfying

$$u_1(1, t) = u_2(0, t) = \dots = u_k(0, t) \quad (30)$$

$$c_2 u_{2,x}(0, t) + \dots + c_k u_{k,x}(0, t) - c_1 u_{1,x}(1, t) = -\alpha u_{1,t}(1, t) \quad (31)$$

Introduce the Riemann invariants

$$d_i(x, t) := u_{i,t}(x, t) - c_i u_{i,x}(x, t), \quad (32)$$

$$s_i(x, t) := u_{i,t}(x, t) + c_i u_{i,x}(x, t) \quad (33)$$

for all  $i \in \mathcal{I}$ . Then the following result holds.

**Lemma 2.2.** 1. If  $\alpha \neq k$ , then  $s_1(1, t), d_2(0, t), \dots, d_k(0, t)$  can be expressed in a unique way as functions of  $d_1(1, t), s_2(0, t), \dots, s_k(0, t)$ . In particular, if  $\alpha = k - 2$ , we obtain

$$s_1(1, t) = \sum_{i=2}^k s_i(0, t). \quad (34)$$

2. If  $\alpha = k$ , then the existence of a solution to (12) and (30)-(31) implies

$$d_1(1, t) + \sum_{i=2}^k s_i(0, t) = 0. \quad (35)$$

As a consequence, the initial data of a smooth solution should satisfy the compatibility condition

$$v_1^0(1 - c_1 t) - c_1 u_{1,x}^0(1 - c_1 t) + \sum_{i=2}^k [v_i^0(c_i t) + c_i u_{i,x}^0(c_i t)] = 0$$

$$\text{for } 0 \leq t \leq \min_{1 \leq i \leq k} c_i^{-1}. \quad (36)$$

*Proof of Lemma 2.2.* Using Riemann invariants, we see that (12) and (30)-(31) are transformed into

$$d_{i,t} + c_i d_{i,x} = 0, \quad i = 1, \dots, k, \quad (37)$$

$$s_{i,t} - c_i s_{i,x} = 0, \quad i = 1, \dots, k, \quad (38)$$

$$s_1(1, t) + d_1(1, t) = s_2(0, t) + d_2(0, t) = \dots = s_k(0, t) + d_k(0, t), \quad (39)$$

$$\sum_{i=2}^k [s_i(0, t) - d_i(0, t)] - (s_1(1, t) - d_1(1, t)) = -\alpha(s_1(1, t) + d_1(1, t)). \quad (40)$$

To simplify the notations, we write  $s_1$  for  $s_1(1, t)$ ,  $s_2$  for  $s_2(0, t)$ , etc. Then (39)-(40) can be written

$$s_1 + d_1 = d_i + s_i, \quad i = 2, \dots, k, \quad (41)$$

$$(1 - \alpha)s_1 + d_2 + \dots + d_k = (1 + \alpha)d_1 + s_2 + \dots + s_k \quad (42)$$

We readily infer from (41) that

$$s_1 - d_2 = -d_1 + s_2, \quad (43)$$

$$d_2 - d_3 = -s_2 + s_3, \quad (44)$$

$$\vdots$$

$$d_{k-1} - d_k = -s_{k-1} + s_k. \quad (45)$$

Adding the  $k - 1$  equations in (41) results in

$$(k - 1)s_1 - \sum_{i=2}^k d_i = (1 - k)d_1 + \sum_{i=2}^k s_i$$

Subtracting this last equation from (42), we obtain

$$2 \sum_{i=2}^k d_i = (k + \alpha)d_1 + (k + \alpha - 2)s_1 = 2d_1 + (k + \alpha - 2)(d_1 + s_1).$$



Combined to the relation  $d_1 + s_1 = d_k + s_k$ , this yields

$$\sum_{i=2}^k d_i = d_1 + \left(\frac{k+\alpha}{2} - 1\right)(d_k + s_k).$$

Using this relation in (42) together with the relation  $s_1 = d_k + s_k - d_1$ , we obtain

$$(k-\alpha)d_k = 2d_1 + 2\sum_{i=2}^{k-1} s_i + (\alpha - k + 2)s_k. \quad (46)$$

Thus, if  $\alpha \neq k$ , we infer from (43)-(46) that  $s_1(1, t), d_2(0, t), \dots, d_k(0, t)$  can be expressed in a unique way as functions of  $d_1(1, t), s_2(0, t), \dots, s_k(0, t)$ . In particular, if  $\alpha = k - 2$ , then (46) becomes

$$d_k = d_1 + \sum_{i=2}^{k-1} s_i. \quad (47)$$

Adding (43), (44), ..., (45) and (47) yields (34). Finally, if  $\alpha = k$ , then (46) gives (35). Computing  $d_i$  and  $s_i$  using (52)-(53) (see below), and replacing  $s_i^0$  and  $d_i^0$  by their expressions in terms of  $u_i^0$  and  $v_i^0$ , we obtain (36).  $\square$

Let us proceed to the proof of Theorem 2.1. If (28) is not satisfied, picking some initial data  $(u_i^0, v_i^0)_{i \in \mathcal{I}} \in \mathcal{D}(A_T)$  that does not satisfy the condition

$$v_n^0(1 - c_n t) - c_n u_{n,x}^0(1 - c_n t) + \sum_{i \in \mathcal{I}_n} [v_i^0(c_i t) + c_i u_{i,x}^0(c_i t)] = 0, \quad 0 \leq t \leq \min_{i \in \{n\} \cup \mathcal{I}_n} c_i^{-1} \quad (48)$$

around a node  $n \in \mathcal{N}_M$  for which  $\alpha_n = k_n$  (the existence of such an initial data is obvious, since the conditions (20)-(27) involve only the values of  $u_i$  and  $v_i$  or their derivatives at the nodes), we infer from Lemma 2.2 that system (12)-(16) and (19) does not admit any solution  $(u_i, v_i)_{i \in \mathcal{I}} \in C(\mathbb{R}^+; \mathcal{D}(A_T))$ . This shows that  $A_T$  is not the generator of a continuous semigroup on  $\mathcal{H}$ .

Conversely, assume that (28) is satisfied. We aim to construct by a fixed-point procedure a solution to (12)-(16) and (19). Pick any  $U^0 = (u_i^0, v_i^0)_{i \in \mathcal{I}} \in \mathcal{H}$  and any  $T > \sup_{1 \leq i \leq N} c_i^{-1}$ . Set

$$d_i^0 := v_i^0 - c_i u_{i,x}^0, \quad s_i^0 := v_i^0 + c_i u_{i,x}^0, \quad i = 1, \dots, N.$$

Let  $K := \sum_{n \in \mathcal{N}_M} k_n$ . Pick any number  $\rho \in (0, 1)$ , and introduce the Hilbert space  $\mathcal{E} := L^2_{\rho^t dt}(0, T)^K$  endowed with the norm

$$\|(x_1, x_2, \dots, x_K)\|_{\mathcal{E}} := \left( \sum_{i=1}^K \int_0^T |x_i(t)|^2 \rho^t dt \right)^{\frac{1}{2}}.$$

For  $n \in \mathcal{N}_M$ , let  $m_n := \inf_{j \in \mathcal{I}_n} j$ , and let  $p_1 := 1$  and

$$p_n := 1 + \sum_{m \in \mathcal{N}_M, m < n} k_m, \quad \text{for } n \in \mathcal{N}_M.$$

For instance, with the tree drawn in Figure 1, we have

$$\begin{aligned} \mathcal{N}_M &= \{1, 2, 3, 5, 6, 7\}, \\ (k_1, k_2, k_3, k_5, k_6, k_7) &= (3, 3, 2, 4, 3, 3), \end{aligned}$$

$$\begin{aligned} (m_1, m_2, m_3, m_5, m_6, m_7) &= (2, 4, 6, 7, 10, 12), \\ \text{and } (p_1, p_2, p_3, p_5, p_6, p_7) &= (1, 4, 7, 9, 13, 16). \end{aligned}$$

Let

$$X(t) := (x_1(t), \dots, x_K(t)) = (\dots, d_n(1, t), s_{m_n}(0, t), \dots, s_{m_n+k_n-2}(0, t), \dots)$$

where  $n$  ranges over  $\mathcal{N}_M$ . Note that  $x_1(t) = d_1(1, t)$  and  $x_K(t) = s_N(0, t)$ . For the tree drawn in Figure 1, we have

$$\begin{aligned} X(t) &= (x_1(t), \dots, x_{18}(t)) \\ &= (d_1(1, t), s_2(0, t), s_3(0, t), d_2(1, t), s_4(0, t), s_5(0, t), d_3(1, t), s_6(0, t), \\ &\quad d_5(1, t), s_7(0, t), s_8(0, t), s_9(0, t), d_6(1, t), s_{10}(0, t), s_{11}(0, t), d_7(1, t), \\ &\quad s_{12}(0, t), s_{13}(0, t)). \end{aligned}$$

Finally, let  $\mathcal{S}$  denote the set of the indices of the components of  $X(t)$  corresponding to edges whose final points are simple nodes different from the root; that is

$$\mathcal{S} := \{n \in \{2, \dots, K\}, \exists q_n \in \mathcal{N}_S, x_n(t) = s_{q_n}(0, t)\}.$$

For the tree in Figure 1, we have  $\mathcal{S} = \{5, 11, 12, 14, 15, 17, 18\}$  with

$$(q_5, q_{11}, q_{12}, q_{14}, q_{15}, q_{17}, q_{18}) = (4, 8, 9, 10, 11, 12, 13).$$

Let

$$\mathcal{E}_0 := \{(x_1, \dots, x_K) \in \mathcal{E}; x_1(t) = 0 \forall t \geq c_1^{-1} \text{ and } x_n(t) = 0 \forall n \in \mathcal{S}, \forall t \geq c_{q_n}^{-1}\}.$$

We define a map  $P : X = (x_1, \dots, x_K) \in \mathcal{E}_0 \rightarrow \tilde{X} = (\tilde{x}_1, \dots, \tilde{x}_K) \in \mathcal{E}_0$  as follows. Pick any  $X = (x_1, \dots, x_K) \in \mathcal{E}_0$  and any  $n \in \mathcal{N}_M$ . By Lemma 2.2, there exists a matrix  $A_n \in \mathbb{R}^{k_n \times k_n}$  such that the Riemann invariants associated with the solution of (12)-(16) and (19) satisfy

$$\begin{pmatrix} s_n(1, t) \\ d_{m_n}(0, t) \\ \vdots \\ d_{m_n+k_n-2}(0, t) \end{pmatrix} = A_n \begin{pmatrix} d_n(1, t) \\ s_{m_n}(0, t) \\ \vdots \\ s_{m_n+k_n-2}(0, t) \end{pmatrix}.$$

Then, we set for all  $t \geq 0$

$$\begin{pmatrix} s_n(1, t) \\ d_{m_n}(0, t) \\ \vdots \\ d_{m_n+k_n-2}(0, t) \end{pmatrix} := A_n \begin{pmatrix} x_{p_n}(t) \\ x_{p_n+1}(t) \\ \vdots \\ x_{p_n+k_n-1}(t) \end{pmatrix}. \tag{49}$$

(Recall that  $x_{p_n}(t)$  stands for  $d_n(1, t)$ .) Next we set

$$d_1(0, t) = 0, \quad t \geq 0, \tag{50}$$

$$s_m(1, t) = 0, \quad m \in \mathcal{N}_S \setminus \{0\}, \quad t \geq 0. \tag{51}$$

Note that  $d_i(0, t)$  and  $s_i(1, t)$  have been defined for all  $i \in \{1, \dots, N\}$  and all  $t \geq 0$ . Next, solving (37)-(38), we set for all  $i \in \{1, \dots, N\}$  and all  $(x, t) \in [0, 1] \times \mathbb{R}^+$

$$d_i(x, t) := \begin{cases} d_i^0(x - c_i t) & \text{if } 0 \leq x - c_i t \leq 1, \\ d_i(0, t - c_i^{-1}x) & \text{if } t > x/c_i, \end{cases} \tag{52}$$

and

$$s_i(x, t) := \begin{cases} s_i^0(x + c_i t) & \text{if } 0 \leq x + c_i t \leq 1, \\ s_i(1, t - c_i^{-1}(1 - x)) & \text{if } t > (1 - x)/c_i. \end{cases} \tag{53}$$

Finally, we set for  $n \in \mathcal{N}_M$  and  $t \geq 0$

$$\begin{pmatrix} \tilde{x}_{p_n}(t) \\ \tilde{x}_{p_n+1}(t) \\ \vdots \\ \tilde{x}_{p_n+k_n-1}(t) \end{pmatrix} := \begin{pmatrix} d_n(1, t) \\ s_{m_n}(0, t) \\ \vdots \\ s_{m_n+k_n-2}(0, t) \end{pmatrix}. \quad (54)$$

and  $P(X) := \tilde{X} = (\tilde{x}_1, \dots, \tilde{x}_K)$ .

From (50)-(53), we have that

$$d_1(1, t) = 0, \quad \text{for } t \geq c_1^{-1}, \quad (55)$$

$$s_m(0, t) = 0, \quad \text{for } m \in \mathcal{N}_S \setminus \{0\}, t \geq c_m^{-1}. \quad (56)$$

It follows that  $P$  is a map from  $\mathcal{E}_0$  into itself. Let us check that it is a contraction for  $\rho$  small enough. Let  $X^1 = (x_1^1, \dots, x_K^1)$  and  $X^2 = (x_1^2, \dots, x_K^2)$  be given in  $\mathcal{E}_0$ . In what follows,  $c$  denotes a constant that may vary from line to line. Then, using (49)-(54), we have that

$$\begin{aligned} \|P(X_1) - P(X_2)\|_{\mathcal{E}}^2 &\leq c \sum_{j=1}^K \sum_{i=1}^N \int_{c_i^{-1}}^T |x_j^1(t - c_i^{-1}) - x_j^2(t - c_i^{-1})|^2 \rho^t dt \quad (57) \\ &\leq c(\max_{i \in \mathcal{I}} \rho^{c_i^{-1}}) \|X^1 - X^2\|_{\mathcal{E}}^2. \quad (58) \end{aligned}$$

This proves that  $P$  is a contraction in  $\mathcal{E}_0$  for  $\rho > 0$  small enough. It follows from the contraction principle that  $P$  has a (unique) fixed-point in  $\mathcal{E}_0$ . It is then easy to check that the Riemann invariants  $d_i, s_i, 1 \leq i \leq N$ , defined along (52)-(53), solve (37)-(38) in the distributional sense and satisfy (39)-(40) almost everywhere. Using again (52)-(53), one has that for any  $i \in \mathcal{I}$

$$s_i(x, 0) = s_i^0(x), \quad d_i(x, 0) = d_i^0(x), \quad \text{for a.e. } x \in [0, 1].$$

We can therefore define for all  $i \in \mathcal{I}$  and all  $T > \sup_{1 \leq i \leq N} c_i^{-1}$  a function  $u_i \in H^1((0, 1) \times (0, T))$  by

$$u_{i,t} = \frac{1}{2}(s_i + d_i) =: v_i, \quad u_{i,x} = \frac{1}{2c_i}(s_i - d_i),$$

the constant of integration being chosen so that

$$u_i(x, t) = u_i^0(x) + \int_0^t v_i(x, s) ds \quad \text{for a.e. } (x, t) \in (0, 1) \times (0, T).$$

Then  $(u_i, v_i) \in C(\mathbb{R}^+, H^1(0, 1) \times L^2(0, 1))$ , and (22) follows from (39). We infer that  $(u_i, v_i)_{i \in \mathcal{I}}$  is a (weak) solution of (12)-(16) and (19) which is continuous in time with values in  $\mathcal{H}$ . Set  $S(t)U^0 := (u_i(t), v_i(t))_{i \in \mathcal{I}}$ . Then it can be seen that  $(S(t))_{t \geq 0}$  is a strongly continuous semigroup in  $\mathcal{H}$  whose generator is  $A_T$ . The proof of Theorem 2.1 is complete.  $\square$

**3. The example of a tree with two internal nodes.** In this section, we consider a particular case which can be treated in a direct way, and which shows the importance of condition (7) in Theorem 1.1.

To prove on a given example that the finite-time stability (to 0 or to constant functions) does not occur, a way consists in finding an eigenvalue of the underlying operator. Indeed, if we can find an eigenvalue, then the corresponding exponential solution does not steer 0 in finite time. If, in addition, the eigenvalue is different from 0, then the corresponding exponential solution does not take a constant value

after some finite time. The explicit computation of the eigenvalues can be achieved when the geometry is sufficiently simple, typically  $d \leq 3$ .

In this section, we assume now  $\mathcal{T}$  is a tree with  $N + 1$  nodes, two of which being multiple ( $d = 3, \mathcal{N}_M = \{1, 2\}, k_1 \geq 2, k_2 \geq 2, k_1 + k_2 = N + 1$ ), and we consider the boundary conditions (14)-(16) and (19). (See Figure 2.) We will let  $\alpha_1$  and

● Transparent boundary condition




FIGURE 2. A bone-shaped tree with  $k_1 = k_2 = 3$ .

$\alpha_2$  range over  $\mathbb{R}$ , assuming only that (28) holds. In particular, when  $\alpha_1 = \alpha_2 = 0$ , there is no damping at the internal nodes  $n = 1, 2$ . We shall show that the finite-time stabilization cannot hold in that case, because of the (partial but continuous) bounces of waves at the internal nodes. Note that for this geometry, condition (7) reads

$$\alpha_1 = k_1 - 2, \quad \alpha_2 = k_2 - 2. \tag{59}$$

Here, we shall show that there is an eigenvalue (and actually a sequence of eigenvalues, so that the finite-time stability to constant functions fails) if, and only if, both  $\alpha_1 \neq k_1 - 2$  and  $\alpha_2 \neq k_2 - 2$ . Notice that this condition is stronger than  $(\alpha_1, \alpha_2) \neq (k_1 - 2, k_2 - 2)$ . We shall also prove that, when

$$(\alpha_1, \alpha_2) \in \{k_1 - 2\} \times (\mathbb{R} \setminus \{k_2\}) \cup (\mathbb{R} \setminus \{k_1\}) \times \{k_2 - 2\}, \tag{60}$$

then the finite-time stability (to constant functions) occurs. We conclude that, when  $d = 3$  and transparent boundary conditions are imposed at all the external nodes, a necessary and sufficient condition for the finite-time stability (to constant functions) is (60). The interpretation is that the nodes satisfying (7) and for which all the adjacent nodes but one are external, are “transparent” and can be “removed” from the tree.

Let  $(\alpha_1, \alpha_2) \in \mathbb{R}^2$  be given. The operator  $A_T$  reads then

$$A_T((u_i, v_i)_{i \in \mathcal{I}}) = (v_i, c_i^2 u_i'')_{i \in \mathcal{I}}$$

with domain

$$\begin{aligned} D(A_T) = \{ & (u_i, v_i)_{i \in \mathcal{I}} \in \mathcal{H}; (u_i, v_i)_{i \in \mathcal{I}} \in \prod_{i \in \mathcal{I}} [H^2(0, 1) \times H^1(0, 1)], \\ & c_1 u_1'(0) = v_1(0), \quad c_i u_i'(1) = -v_i(1) \text{ for } i \in \{3, \dots, N\}, \\ & \sum_{2 \leq i \leq k_1} c_i u_i'(0) - c_1 u_1'(1) = -\alpha_1 v_1(1), \\ & \sum_{k_1+1 \leq i \leq N} c_i u_i'(0) - c_2 u_2'(1) = -\alpha_2 v_2(1), \end{aligned}$$

$$\begin{aligned}(u_i(0), v_i(0)) &= (u_1(1), v_1(1)) \quad \text{for } 2 \leq i \leq k_1, \\ (u_i(0), v_i(0)) &= (u_2(1), v_2(1)) \quad \text{for } k_1 + 1 \leq i \leq N.\end{aligned}$$

where  $' = d/dx$ ,  $'' = d^2/dx^2$ , etc.

Setting  $U = (u_i, v_i)_{i \in \mathcal{I}}$ , we see that (12)-(16) and (19) may be written as

$$U_t = A_T U, \quad (61)$$

$$U(0) = U_0 = (u_i^0, u_i^1)_{i \in \mathcal{I}}. \quad (62)$$

If  $A_T U_0 = \lambda U_0$  with  $U_0 \neq 0$  and  $\lambda \neq 0$ , then the solution  $U$  of (61)-(62) reads  $U(t) = e^{\lambda t} U_0$  (exponential solution), and then the finite-time stabilization to constant functions cannot hold.

**Proposition 1.** *Let  $\mathcal{T}$  denote a tree with  $N$  edges and two internal nodes ( $\mathcal{N}_M = \{1, 2\}$ ), and assume that*

$$\alpha_1 \neq k_1 \text{ and } \alpha_2 \neq k_2. \quad (63)$$

*Then the operator  $A_T$  has at least one eigenvalue if, and only if,*

$$\alpha_1 \neq k_1 - 2 \text{ and } \alpha_2 \neq k_2 - 2. \quad (64)$$

*Furthermore, if (64) holds, then the discrete spectrum of  $A_T$  is  $\sigma_d(A_T) = \{\lambda_k; k \in \mathbb{Z}\}$  where*

$$\lambda_k = \frac{c_2}{2} \log_{-\frac{\pi}{2}} \left( \frac{(2 + \alpha_1 - k_1)(2 + \alpha_2 - k_2)}{(\alpha_1 - k_1)(\alpha_2 - k_2)} \right) + ic_2 k \pi. \quad (65)$$

*In particular, the finite-time stability to constant functions does not hold for (12)-(16) and (19). Finally, if (60) is satisfied, then the finite-time stability to constant functions holds.*

In (65),  $\log_{-\frac{\pi}{2}}$  denotes the usual determination of the logarithm in  $\mathbb{C} \setminus i\mathbb{R}^-$ . Thus

$$\log_{-\frac{\pi}{2}}(z) = \begin{cases} \log |z| & \text{if } z \in (0, +\infty), \\ \log |z| + i\pi & \text{if } z \in (-\infty, 0). \end{cases}$$

*Proof.* First,  $A_T$  generates a strongly continuous semigroup of operators in  $\mathcal{H}$  by (63) and Theorem 2.1. Let  $\lambda \in \mathbb{C}$  and  $U = (u_i, v_i)_{i \in \mathcal{I}} \in D(A_T)$ . Then the equation  $A_T U = \lambda U$  is equivalent to the following system

$$(v_i, c_i^2 u_i'') = \lambda(u_i, v_i) \quad (66)$$

$$c_1 u_1'(0) = v_1(0) \quad (67)$$

$$c_i u_i'(1) = -v_i(1), \quad 3 \leq i \leq N \quad (68)$$

$$\sum_{2 \leq i \leq k_1} c_i u_i'(0) - c_1 u_1'(1) = -\alpha_1 v_1(1) \quad (69)$$

$$\sum_{k_1+1 \leq i \leq N} c_i u_i'(0) - c_2 u_2'(1) = -\alpha_2 v_2(1) \quad (70)$$

$$u_i(0) = u_1(1), \quad 2 \leq i \leq k_1, \quad (71)$$

$$u_i(0) = u_2(1), \quad k_1 + 1 \leq i \leq N. \quad (72)$$

Note that the conditions  $v_i(0) = v_1(1)$  for  $2 \leq i \leq k_1$  and  $v_i(0) = v_2(1)$  for  $k_1 + 1 \leq i \leq N$  are satisfied whenever (66) and (71)-(72) hold. (66) is easily solved as

$$u_i(x) = a_i e^{\lambda x/c_i} + b_i e^{-\lambda x/c_i}, \quad v_i = \lambda u_i, \quad i \in \mathcal{I}, \quad (73)$$

where  $a_i, b_i \in \mathbb{C}$  are constants to be determined. Substituting the above expression of  $u_i(x)$  in (67)-(72) yields the system

$$\lambda b_1 = 0, \quad (74)$$

$$\lambda a_i = 0, \quad 3 \leq i \leq N, \quad (75)$$

$$\lambda \sum_{2 \leq i \leq k_1} (a_i - b_i) - \lambda(a_1 e^{\lambda/c_1} - b_1 e^{-\lambda/c_1}) = -\alpha_1 \lambda(a_1 e^{\lambda/c_1} + b_1 e^{-\lambda/c_1}), \quad (76)$$

$$\lambda \sum_{k_1+1 \leq i \leq N} (a_i - b_i) - \lambda(a_2 e^{\lambda/c_2} - b_2 e^{-\lambda/c_2}) = -\alpha_2 \lambda(a_2 e^{\lambda/c_2} + b_2 e^{-\lambda/c_2}), \quad (77)$$

$$a_i + b_i = a_1 e^{\lambda/c_1} + b_1 e^{-\lambda/c_1}, \quad 2 \leq i \leq k_1, \quad (78)$$

$$a_i + b_i = a_2 e^{\lambda/c_2} + b_2 e^{-\lambda/c_2}, \quad k_1 + 1 \leq i \leq N. \quad (79)$$

If  $\lambda = 0$ , we infer from (78)-(79) and (73) that  $u_i(x) = a_1 + b_1$  for all  $i \in \mathcal{I}$ , i.e.  $U = \text{const}$ , which is excluded. Assume from now on that  $\lambda \neq 0$ . Then (74)-(79) is equivalent to the system

$$b_1 = 0, \quad (80)$$

$$a_i = 0, \quad 3 \leq i \leq N, \quad (81)$$

$$b_2 = a_1 e^{\lambda/c_1} - a_2, \quad (82)$$

$$b_i = a_1 e^{\lambda/c_1}, \quad 3 \leq i \leq k_1, \quad (83)$$

$$b_i = a_2 e^{\lambda/c_2} + b_2 e^{-\lambda/c_2}, \quad k_1 + 1 \leq i \leq N, \quad (84)$$

$$2a_2 + (\alpha_1 - k_1)e^{\lambda/c_1} a_1 = 0, \quad (85)$$

$$\begin{aligned} & [(-N + k_1 - 1 + \alpha_2)e^{\lambda/c_2} + (N - k_1 - 1 - \alpha_2)e^{-\lambda/c_2}]a_2 \\ & + (-N + k_1 + 1 + \alpha_2)e^{-\lambda/c_2} e^{\lambda/c_1} a_1 = 0. \end{aligned} \quad (86)$$

The existence of a nontrivial solution  $((a_1, a_2) \neq (0, 0))$  holds if, and only if, the determinant of the system (85)-(86) in  $e^{\lambda/c_1} a_1$  and  $a_2$  vanishes, i.e.

$$(2 + \alpha_1 - k_1)(-N + k_1 + 1 + \alpha_2)e^{-\lambda/c_2} - (\alpha_1 - k_1)(-N + k_1 - 1 + \alpha_2)e^{\lambda/c_2} = 0.$$

Since  $-N + k_1 = 1 - k_2$ , this can be expressed as

$$(2 + \alpha_1 - k_1)(2 + \alpha_2 - k_2)e^{-\lambda/c_2} - (\alpha_1 - k_1)(\alpha_2 - k_2)e^{\lambda/c_2} = 0.$$

Using (63), the last equation is equivalent to

$$e^{\frac{2\lambda}{c_2}} = \frac{(2 + \alpha_1 - k_1)(2 + \alpha_2 - k_2)}{(\alpha_1 - k_1)(\alpha_2 - k_2)}. \quad (87)$$

(87) has a solution  $\lambda \in \mathbb{C}$  if and only if  $(2 + \alpha_1 - k_1)(2 + \alpha_2 - k_2) \neq 0$ , and in that case the solutions of (87) read

$$\lambda_k = \frac{c_2}{2} \log_{-\frac{\pi}{2}} \left( \frac{(2 + \alpha_1 - k_1)(2 + \alpha_2 - k_2)}{(\alpha_1 - k_1)(\alpha_2 - k_2)} \right) + ic_2 k \pi, \quad k \in \mathbb{Z}.$$

Assume finally that (60) holds, e.g.  $\alpha_1 = k_1 - 2$  and  $\alpha_2 \in \mathbb{R} \setminus \{k_2\}$ . Since transparent boundary conditions are applied at all the external nodes, we have

$$\begin{aligned} s_i(1, t) &= 0, & i = 3, \dots, N, \quad t \geq 0, \\ d_1(0, t) &= 0, & t \geq 0. \end{aligned}$$

This implies

$$s_i(x, t) = 0, \quad i = 3, \dots, N, \quad x \in [0, 1], \quad t \geq c_i^{-1}, \quad (88)$$

$$d_1(x, t) = 0, \quad x \in [0, 1], \quad t \geq c_1^{-1}. \quad (89)$$

It follows from (34) and (88) that

$$s_2(0, t) = s_1(1, t), \quad t \geq \max_{3 \leq i \leq k_1} c_i^{-1}.$$

Combined with the continuity condition  $u_1(1, t) = u_2(0, t)$ , this yields

$$d_2(0, t) = d_1(1, t) = 0, \quad t \geq \max_{i \in \{1\} \cup [3, k_1]} c_i^{-1}.$$

We infer then from Lemma 2.2 (see (43)-(45) and (46)) that  $s_2(1, t) = d_{k_1+1}(0, t) = \dots = d_N(0, t) = 0$  for  $t$  large enough. This in turn implies  $s_1(1, t) = d_3(0, t) = \dots = d_{k_1}(0, t) = 0$  for  $t$  large enough. We conclude that for some constants  $C$  and  $T$ ,  $u_i(x, t) = C$  for all  $i \in \mathcal{I}$ , all  $x \in [0, 1]$  and all  $t \geq T$ .  $\square$

**4. Finite-time extinction.** Pick any tree of depth  $d \geq 1$ , and define the sequence  $(t_i)_{i \in \mathcal{I}}$  as follows

$$t_i = c_i^{-1} \quad \text{if } i \in \mathcal{N}_S \setminus \{0\}, \quad (90)$$

$$t_i = c_i^{-1} + \max_{j \in \mathcal{I}_i} t_j \quad \text{if } i \in \mathcal{N}_M. \quad (91)$$

Set  $T(\mathcal{R}) = t_1$ . Then it is easily seen that  $T(\mathcal{R})$  is the maximum of the quantities

$$c_{i_1}^{-1} + c_{i_2}^{-1} + \dots + c_{i_p}^{-1},$$

where  $p \geq 1$ ,  $i_1 = 1$ ,  $i_{q+1} \in I_{i_q}$  for  $1 \leq q \leq p-1$ , and the final point of the edge of index  $i_q$  is an external node (different from  $\mathcal{R}$ ). Define  $T(\mathcal{T})$  as the largest of the  $T(\mathcal{R})$ 's when the root  $\mathcal{R}$  ranges over  $\mathcal{N}_S$ ; that is, we take as root of the tree any external node, change the numbering of the edges and nodes, and define the corresponding sequences  $(\mathcal{I}_i)_{i \in \mathcal{I}}$  and  $(t_i)_{i \in \mathcal{I}}$ . Obviously,  $T(\mathcal{R}) \leq T(\mathcal{T}) \leq 2T(\mathcal{R})$ .

**Example.** Consider again the tree drawn in Figure 1, and assume for simplicity that  $c_i = 1$  for all  $i \in [1, 13]$ . Then  $T(\mathcal{R}) = 5$  and  $T(\mathcal{T}) = 7$ . Indeed, if we take the node of index  $n = 12$  as (new) root, we obtain  $T(\mathcal{R}_{n=12}) = 7$ . Similarly, we see that  $T(\mathcal{R}_{n=13}) = 7$ ,  $T(\mathcal{R}_{n=8}) = T(\mathcal{R}_{n=9}) = 6$ ,  $T(\mathcal{R}_{n=4}) = 5$ , and  $T(\mathcal{R}_{n=10}) = T(\mathcal{R}_{n=11}) = 7$ .

**Theorem 4.1.** Let  $\mathcal{T}$  be a tree of root  $\mathcal{R}$ , and let  $T(\mathcal{R})$  and  $T(\mathcal{T})$  be as above. Assume that the sequence  $(\alpha_n)_{n \in \mathcal{N}_M}$  satisfies the condition

$$\alpha_n = k_n - 2 \quad n \in \mathcal{N}_M. \quad (92)$$

Pick any initial data  $U_0 = \{(u_i^0, u_i^1)\}_{i \in \mathcal{I}} \in \mathcal{H}$ .

(i) If  $U_0 \in \mathcal{H}_0$ , then the solution  $(u_i)_{i \in \mathcal{I}}$  of (12)-(16) and (17) satisfies

$$u_i(\cdot, t) \equiv 0, \quad \forall t \geq 2T(\mathcal{R}), \quad \forall i \in \mathcal{I}; \quad (93)$$

(ii) The solution  $(u_i)_{i \in \mathcal{I}}$  of (12)-(16) and (18) satisfies for some number  $C \in \mathbb{R}$

$$u_i(\cdot, t) \equiv C, \quad \forall t \geq 2T(\mathcal{R}), \quad \forall i \in \mathcal{I}. \quad (94)$$

(iii) The solution  $(u_i)_{i \in \mathcal{I}}$  of (12)-(16) and (19) satisfies for some number  $C \in \mathbb{R}$

$$u_i(\cdot, t) \equiv C, \quad \forall t \geq T(\mathcal{T}), \quad \forall i \in \mathcal{I}. \quad (95)$$

**Remark 2.** It is likely that the extinction time  $T_e$  (i.e. the least time after which solutions remain constant) is given by  $2T(\mathcal{R})$  in the cases (i) and (ii), and  $T(\mathcal{T})$  in case (iii), so that the above results are sharp. Actually, for one string, it is well known that  $T_e = 2/c_1$  for the solutions of (12)-(16) and (17) (or for the solutions of (12)-(16) and (18)), while  $T_e = 1/c_1$  for the solutions of (12)-(16) and (19) (see e.g. [12, 14, 17] and the references therein).

*Proof.* We use again the Riemann invariants  $d_i, s_i$  defined in (32)-(33) that satisfy the transport equations (37)-(38). We need the following

**Lemma 4.2.** *Let  $\mathcal{T}$  be a tree, and let the sequence  $(t_i)_{i \in \mathcal{I}}$  be as in (90)-(91). Assume that the sequence  $(\alpha_n)_{n \in \mathcal{N}_M}$  satisfies (92). Then for any  $U_0 \in \mathcal{H}$  and any solution  $(u_i)_{i \in \mathcal{I}}$  of (12)-(16), with corresponding Riemann invariants  $d_i, s_i$ , we have for all  $i \in \mathcal{I}$*

$$s_i(x, t) = 0 \quad \forall x \in [0, 1], \forall t \geq t_i. \tag{96}$$

*Proof of Lemma 4.2.* We argue by induction on the depth  $d$  of the tree. If  $d = 1$ , then there is only one edge ( $\mathcal{I} = \{1\}$ ) and  $s_1$  solves

$$s_{1,t} - c_1 s_{1,x} = 0, \quad t > 0, \quad 0 < x < 1, \tag{97}$$

$$s_1(1, t) = 0, \quad t > 0, \tag{98}$$

$$s_1(\cdot, 0) = s_1^0 := v_1^0 + c_1 u_{1,x}^0. \tag{99}$$

Then it is easily seen that

$$s_1(x, t) = \begin{cases} s_1^0(x + c_1 t) & \text{if } x + c_1 t \leq 1, \\ 0 & \text{if } x + c_1 t \geq 1. \end{cases} \tag{100}$$

Thus

$$s_1(x, t) = 0 \quad \forall x \in [0, 1], \forall t \geq c_1^{-1}$$

and (96) is established for  $d = 1$ .

Assume now Lemma 4.2 established for any tree of depth at most  $d - 1$ , where  $d \geq 2$ . Pick a tree  $\mathcal{T}$  of depth  $d$ , and a sequence  $(\alpha_n)_{n \in \mathcal{N}_M}$  satisfying  $(\mathcal{P})$ . Denote by  $\mathcal{R}'$  the node of index  $n = 1$ , and by  $\mathcal{T}_i$ , for  $i = 2, \dots, k_1$ , the subtree of  $\mathcal{T}$  of root  $\mathcal{R}'$  and of first edge the edge of  $\mathcal{T}$  of index  $i$ . Since  $\mathcal{T}_i$  is of depth at most  $d - 1$ , we infer from the induction hypothesis that for  $i > 1$

$$s_i(x, t) = 0 \quad \forall x \in [0, 1], \forall t \geq t_i. \tag{101}$$

It remains to prove (96) for  $i = 1$ . Since the condition (92) is satisfied for  $n = 1$ , we infer from (34) that

$$s_1(1, t) = \sum_{i=2}^{k_1} s_i(0, t), \quad \forall t \geq 0.$$

It follows then from (101) that

$$s_1(1, t) = 0 \quad \forall t \geq \max_{i \in \mathcal{I}_1} t_i.$$

Finally, using (97), we infer that

$$s_1(x, t) = 0 \quad \forall x \in [0, 1], \forall t \geq c_1^{-1} + \max_{i \in \mathcal{I}_1} t_i = t_1.$$

The proof of Lemma 4.2 is complete. □

Let us go back to the proof of Theorem 4.1.



(i) Assume first that  $U_0 \in \mathcal{H}_0$ , and let  $(u_i)_{i \in \mathcal{I}}$  denote the solution of (12)-(16) and (17). From Lemma 4.2, we have that for all  $i \in \mathcal{I}$

$$s_i(x, t) = 0 \quad \forall x \in [0, 1], \forall t \geq T(\mathcal{R}). \quad (102)$$

From (17), we infer that  $d_1(0, t) + s_1(0, t) = 0$  for all  $t \geq 0$ , and hence

$$d_1(0, t) = 0, \quad \forall t \geq T(\mathcal{R}).$$

Using (37), we infer that

$$d_1(x, t) = 0, \quad \forall x \in [0, 1], \forall t \geq c_1^{-1} + T(\mathcal{R}).$$

Combined with (43)-(45) (with  $k = k_1$ ) and (102), this yields

$$d_2(0, t) = \dots = d_{k_1}(0, t) = 0, \quad \forall t \geq c_1^{-1} + \max_{i \in \mathcal{I}_1} c_i^{-1} + T(\mathcal{R}).$$

Using the second definition of  $T(\mathcal{R})$  and proceeding inductively, we arrive to

$$d_i(x, t) = 0 \quad \forall i \in \mathcal{I}, \forall x \in [0, 1], \forall t \geq 2T(\mathcal{R}). \quad (103)$$

Gathering together (102) and (103), we infer the existence of some constant  $C \in \mathbb{R}$  such that

$$u_i(x, t) = C, \quad \forall i \in \mathcal{I}, \forall x \in [0, 1], \forall t \geq 2T(\mathcal{R}).$$

Using (17), we see that  $C = 0$ . This proves that solutions of (12)-(16) and (17) are null for  $t \geq T(\mathcal{R})$ . Combined with the strong continuity of the semigroup  $(e^{tA_D})_{t \geq 0}$  in  $\mathcal{H}_0$ , this yields the finite-time stability.

(ii) Assume now that  $u_0 \in \mathcal{H}$  and let  $(u_i)_{i \in \mathcal{I}}$  denote the solution of (12)-(16) and (18). From (17), we infer that  $d_1(0, t) - s_1(0, t) = 0$  for all  $t \geq 0$ . The same proof as in (i) then yields

$$s_i(x, t) = d_i(x, t) = 0, \quad \forall i \in \mathcal{I}, \forall x \in [0, 1], \forall t \geq 2T(\mathcal{R}).$$

Thus there exists a constant  $C \in \mathbb{R}$  such that

$$u_i(x, t) = C, \quad \forall i \in \mathcal{I}, \forall x \in [0, 1], \forall t \geq 2T(\mathcal{R}).$$

(iii) Pick a solution  $(u_i)_{i \in \mathcal{I}}$  of (12)-(16) and (19). Then it follows from Lemma 4.2 that for all  $i \in \mathcal{I}$

$$s_i(x, t) = 0 \quad \forall x \in [0, 1], \forall t \geq T(\mathcal{R}). \quad (104)$$

For any given  $i \in \mathcal{I}$ , we pick a sequence  $i_1 < i_2 < \dots < i_p$  such that  $i_1 = 1$ ,  $i = i_q$  for some  $q \in [1, p]$ , and the final point of the edge of index  $i_p$  is an external point, that we call  $\tilde{\mathcal{R}}$ . If we exchange  $\mathcal{R}$  and  $\tilde{\mathcal{R}}$ , we notice that  $d_i$  is linked to the  $\tilde{s}_j$ 's (associated with the new root  $\tilde{\mathcal{R}}$ ) by:

$$d_i(x, t) = \tilde{s}_{i_p - i + 1}(1 - x, t).$$

We infer that

$$d_i(x, t) = 0 \quad \forall x \in [0, 1], \forall t \geq T(\mathcal{T}). \quad (105)$$

Therefore, there exists a constant  $C \in \mathbb{R}$  such that

$$u_i(x, t) = C, \quad \forall i \in \mathcal{I}, \forall x \in [0, 1], \forall t \geq T(\mathcal{T}).$$

The proof of Theorem 4.1 is complete.  $\square$

5. **Sharpness of the condition (92).** The condition (92), which is sufficient to yield the finite-time stability, is expected to be also necessary in some situations. (However, we already noticed that it was not necessary for a tree with two internal nodes.)

Here we consider a star-shaped tree, with the homogeneous Dirichlet boundary condition at one external node and the transparent boundary conditions at the other external nodes. We consider any possible value of the coefficient  $\alpha$  at the internal node, and exhibit an eigenvalue of the underlying operator when (28) holds and (92) fails.

Assume that  $\mathcal{T}$  is a star-shaped tree with  $N$  edges ( $d = 2, k_1 = N$ ), and consider the boundary conditions (14)-(16) and (17). (See Figure 3.)




FIGURE 3. A star-shaped tree.

We assume that  $\alpha_1 \neq N$ , so that the system (12)-(16) and (17) is well-posed in  $\mathcal{H}_0$  according to Theorem 2.1. According to Theorem 4.1, there is a finite-time stabilization when  $\alpha_1 = N - 2$ . We shall show that this condition is sharp, i.e. that a finite-time stabilization cannot hold if  $\alpha_1 \notin \{N - 2, N\}$ .

Let  $\alpha_1 \in \mathbb{R}$  be given. The operator  $A_D$  reads

$$A_D((u_i, v_i)_{i \in \mathcal{I}}) = (v_i, c_i^2 u_i'')_{i \in \mathcal{I}}$$

with

$$D(A_D) = \{(u_i, v_i)_{i \in \mathcal{I}} \in \mathcal{H}_0; (v_i, c_i^2 u_i'')_{i \in \mathcal{I}} \in \mathcal{H}_0, \\ c_i u_i'(1) = -v_i(1) \text{ for } 2 \leq i \leq N, \sum_{2 \leq i \leq N} c_i u_i'(0) - c_1 u_1'(1) = -\alpha_1 v_1(1), \\ \text{and } (u_i(0), v_i(0)) = (u_1(1), v_1(1)) \text{ for } 2 \leq i \leq N\},$$

where  $' = d/dx$ ,  $'' = d^2/dx^2$ , etc. Setting  $U := (u_i, v_i)_{i \in \mathcal{I}}$ , we see that (12)-(16) and (17) may be written as

$$U_t = A_D U \tag{106}$$

$$U(0) = U_0 = (u_i^0, u_i^1)_{i \in \mathcal{I}} \tag{107}$$

If  $A_D U_0 = \lambda U_0$  with  $U_0 \neq 0$ , then the solution  $U$  of (106)-(107) reads  $U(t) = e^{\lambda t} U_0$  (exponential solution), and hence  $\|U(t)\|_{\mathcal{H}} = e^{(\text{Re } \lambda)t} \|U_0\|_{\mathcal{H}} > 0$  for all  $t \geq 0$ . Thus if the operator  $A_D$  has at least one eigenvalue, then the finite-time stabilization cannot hold.

**Proposition 2.** Let  $\mathcal{T}$  denote a star-shaped tree with  $N$  edges, and assume that  $\alpha_1 \neq N$ . Then the operator  $A_D$  has at least one eigenvalue if, and only if,

$$\alpha_1 \neq N - 2. \quad (108)$$

Furthermore, if (108) holds, then the discrete spectrum of  $A_D$  is  $\sigma_d(A_D) = \{\lambda_k; k \in \mathbb{Z}\}$  where

$$\lambda_k = \frac{c_1}{2} \log_{-\frac{\pi}{2}} \left( \frac{N-2-\alpha_1}{N-\alpha_1} \right) + ic_1 k \pi \quad (109)$$

In particular, if (108) holds, then the finite-time stabilization of (12)-(16) and (17) in  $\mathcal{H}_0$  fails.

**Remark 3.** (1) It follows from the proof of Proposition 2 that the operator  $A_D$  has no eigenvalue when  $\alpha_1 = N$ . However, as the well-posedness of the system fails by Theorem 2.1, that value of  $\alpha_1$  is not considered for the analysis of the finite-time stability of the system. Actually, there may exist solutions that do not reach a constant value in finite time (pick  $h(t) = t$  in Remark 1 (2)).

(2) If we replace the Dirichlet boundary condition  $u_1(0, t) = 0$  by the transparent boundary condition  $u_{1,t}(0, t) = c_1 u_{1,x}(0, t)$  and take any value  $\alpha_1 \neq N$ , then since  $d_1(0, t) = s_2(1, t) = \dots = s_N(1, t) = 0$  for all  $t \geq 0$ , we infer from (43)-(45) and (46) that  $s_1(1, t) = d_2(0, t) = \dots = d_N(0, t) = 0$  for all  $t \geq \max_{1 \leq i \leq N} c_i^{-1}$ , so that for some constant  $C \in \mathbb{R}$

$$u_i(x, t) = C, \quad \forall i \in [1, N], \forall x \in [0, 1], \forall t \geq 2 \max_{1 \leq i \leq N} c_i^{-1}.$$

*Proof.* Let  $\lambda \in \mathbb{C}$  and  $U = (u_i, v_i)_{i \in \mathcal{I}} \in D(A_D)$ . Then the equation  $A_D U = \lambda U$  is equivalent to the following system

$$(v_i, c_i^2 u_i'') = \lambda(u_i, v_i), \quad 1 \leq i \leq N, \quad (110)$$

$$u_1(0) = 0, \quad (111)$$

$$c_i u_i'(1) = -v_i(1), \quad 2 \leq i \leq N, \quad (112)$$

$$\sum_{2 \leq i \leq N} c_i u_i'(0) - c_1 u_1'(1) = -\alpha_1 v_1(1), \quad (113)$$

$$u_i(0) = u_1(1), \quad 2 \leq i \leq N. \quad (114)$$

Note that the conditions  $v_1(0) = 0$  and  $v_i(0) = v_1(1)$  for  $2 \leq i \leq N$  are satisfied whenever (110)-(111) and (114) hold. (110) is easily solved as

$$u_i(x) = a_i e^{\lambda x/c_i} + b_i e^{-\lambda x/c_i}, \quad v_i(x) = \lambda u_i(x), \quad 1 \leq i \leq N, \quad (115)$$

where  $a_i, b_i \in \mathbb{C}$  are constants to be determined. Substituting the above expression of  $u_i(x)$  in (111)-(114) yields the system

$$a_1 + b_1 = 0, \quad (116)$$

$$\lambda a_i = 0, \quad 2 \leq i \leq N, \quad (117)$$

$$\lambda \sum_{2 \leq i \leq N} (a_i - b_i) - \lambda(a_1 e^{\lambda/c_1} - b_1 e^{-\lambda/c_1}) = -\alpha_1 \lambda(a_1 e^{\lambda/c_1} + b_1 e^{-\lambda/c_1}), \quad (118)$$

$$a_i + b_i = a_1 e^{\lambda/c_1} + b_1 e^{-\lambda/c_1}, \quad 2 \leq i \leq N. \quad (119)$$

If  $\lambda = 0$ , we infer from (115)-(116) and (119) that  $U = 0$ , which is excluded. Assume from now on that  $\lambda \neq 0$ . Then the system (116)-(119) is found to be equivalent to

the system

$$b_1 = -a_1, \tag{120}$$

$$a_i = 0, \quad 2 \leq i \leq N, \tag{121}$$

$$\begin{aligned} & -(N-1)a_1(e^{\lambda/c_1} - e^{-\lambda/c_1}) - a_1(e^{\lambda/c_1} + e^{-\lambda/c_1}) \\ & = -\alpha_1 a_1(e^{\lambda/c_1} - e^{-\lambda/c_1}), \end{aligned} \tag{122}$$

$$b_i = a_1(e^{\lambda/c_1} - e^{-\lambda/c_1}), \quad 2 \leq i \leq N. \tag{123}$$

The existence of a nontrivial solution ( $a_1 \neq 0$ ) holds if, and only if, the coefficient above  $a_1$  in (122) vanishes, i.e.

$$(-N + \alpha_1)e^{\lambda/c_1} + (N - 2 - \alpha_1)e^{-\lambda/c_1} = 0. \tag{124}$$

For  $\alpha_1 \neq N$ , (124) is equivalent to

$$e^{\frac{2\lambda}{c_1}} = \frac{N - 2 - \alpha_1}{N - \alpha_1}.$$

(5) has a solution  $\lambda \in \mathbb{C}$  if and only if  $\alpha_1 \neq N - 2$ , and in that case the solutions of (5) read

$$\lambda_k = \frac{c_1}{2} \log_{-\frac{\pi}{2}} \left( \frac{N - 2 - \alpha_1}{N - \alpha_1} \right) + ic_1 k \pi, \quad k \in \mathbb{Z}. \tag{125}$$

□

**Remark 4.** For  $k \in \mathbb{Z}$  and  $\lambda_k$  as in (125), we introduce the sequence of eigenfunctions  $U_k = ((u_{i,k}, v_{i,k}))_{1 \leq i \leq N, k \in \mathbb{Z}}$  where

$$\begin{aligned} u_{1,k}(x) &= e^{\lambda_k x/c_1} - e^{-\lambda_k x/c_1}, & v_{1,k}(x) &= \lambda_k u_{1,k}(x), \\ u_{i,k}(x) &= (e^{\lambda_k/c_1} - e^{-\lambda_k/c_1})e^{-\lambda_k x/c_1}, & v_{i,k}(x) &= \lambda_k u_{i,k}(x), \quad \text{for } 2 \leq i \leq N. \end{aligned}$$

Then the family  $(a_k U_k)_{k \in \mathbb{Z}}$  may fail to be a Riesz basis in  $\mathcal{H}_0$  for any choice of the sequence of numbers  $(a_k)_{k \in \mathbb{Z}}$ . Consider e.g.  $N = 2$  and  $c_2 = c_1/2$ . Then, for  $N - 2 < \alpha_1 < N$ ,

$$u_{2,k}(x) = (e^{\lambda_k/c_1} - e^{-\lambda_k/c_1})e^{-\log|\frac{N-2-\alpha_1}{N-\alpha_1}|x - i\pi x} e^{-i2k\pi x}.$$

Let  $U = (u_i, v_i)_{i=1,2} \in \mathcal{H}_0$  be given. If  $(a_k U_k)_{k \in \mathbb{Z}}$  is a Riesz basis in  $\mathcal{H}_0$ , then  $U$  can be expanded in terms of the  $U_k$ 's in  $\mathcal{H}_0$  as

$$(u_i, v_i) = \sum_{k \in \mathbb{Z}} d_k a_k (u_{i,k}, v_{i,k}), \quad i = 1, 2$$

for some sequence  $(d_k)_{k \in \mathbb{Z}} \in L^2(\mathbb{Z})$ . Writing

$$e^{\log|\frac{N-2-\alpha_1}{N-\alpha_1}|x + i\pi x} u_2(x) = \sum_{k \in \mathbb{Z}} c_k e^{-i2k\pi x}$$

we have, by harmonicity, that

$$c_k = d_k a_k (e^{\lambda_k/c_1} - e^{-\lambda_k/c_1}), \quad k \in \mathbb{Z},$$

and hence

$$u_1(x) = \sum_{k \in \mathbb{Z}} \frac{c_k}{e^{\lambda_k/c_1} - e^{-\lambda_k/c_1}} (e^{\lambda_k x/c_1} - e^{-\lambda_k x/c_1})$$

in  $L^2(0, 1)$ . Therefore,  $u_1$  is uniquely determined by the  $c_k$ 's, and hence by  $u_2$ , which is a property much stronger than the conditions  $u_1(0) = 0$  and  $u_1(1) = u_2(0)$  present in the definition of  $\mathcal{H}_0$ . This shows that the family  $(a_k U_k)_{k \in \mathbb{Z}}$  is not total in  $\mathcal{H}_0$ .

It is natural to conjecture a decay of all the trajectories like

$$\|U(t)\|_{\mathcal{H}_0} \leq C(\alpha_1) e^{\frac{c_1}{2} \log \left| \frac{N-2-\alpha_1}{N-\alpha_1} \right| t} \|U(0)\|_{\mathcal{H}_0}, \quad t \geq 0, \quad (126)$$

for  $N-2 < \alpha_1 < N$ . (Note that  $\lim_{\alpha_1 \searrow N-2} \log \left| \frac{N-2-\alpha_1}{N-\alpha_1} \right| = -\infty$ .) Without a Riesz basis of eigenvectors in the full space  $\mathcal{H}_0$ , the validity of (126) seems hard to check.

**6. Conclusion.** In this paper, we addressed the issue of the finite-time stabilization of a network of strings. We showed in particular that, when using the homogeneous Dirichlet boundary condition at the root and transparent boundary conditions at the other external nodes, and incorporating in Kirchhoff law the damping term  $\alpha(n)u_t$  with  $\alpha(n) = k(n) - 2$  at each internal node  $n$ , we obtained the finite-time stability of the system. The condition on the coefficients  $\alpha(n)$  proved also to be sharp for a star-shaped tree.

Several questions remain open:

1. Can we obtain a sharp result for a general network? Same question when (12) is replaced by (8)?
2. Can we obtain finite-time stability results for networks containing circuits?

**Acknowledgments.** The authors wish to acknowledge the contribution of anonymous Referees in offering valuable suggestions and in bringing the papers [18, 21] to their attention.

## REFERENCES

- [1] K. Ammari and M. Jellouli, Stabilization of star-shaped networks of strings, *Differential and Integral Equations*, **17** (2004), 1395–1410.
- [2] K. Ammari, M. Jellouli and M. Khenissi, Stabilization of generic trees of strings, *J. Dyn. Control Syst.*, **11** (2005), 177–193.
- [3] A. Bacciotti and L. Rosier, *Liapunov Functions and Stability in Control Theory*, 2<sup>nd</sup> edition, Communications and Control Engineering Series, Springer-Verlag, Berlin, 2005.
- [4] S. P. Bhat and D. S. Bernstein, Finite-time stability of continuous autonomous systems, *SIAM J. Control Optim.*, **38** (2000), 751–766.
- [5] S. Cox and E. Zuazua, The rate at which energy decays in a string damped at one end, *Indiana Univ. Math. J.*, **44** (1995), 545–573.
- [6] R. Dáger and E. Zuazua, Controllability of tree-shaped networks of vibrating strings, *C. R. Acad. Sci. Paris Sér. I Math.*, **332** (2001), 1087–1092.
- [7] R. Dáger and E. Zuazua, *Wave Propagation, Observation and Control in 1-d Flexible Multi-Structures*, Mathématiques & Applications (Berlin) [Mathematics & Applications], 50, Springer-Verlag, Berlin, 2006.
- [8] M. Gugat, Optimal boundary feedback stabilization of a string with moving boundary, *IMA J. Math. Control Inform.*, **25** (2008), 111–121.
- [9] M. Gugat, M. Dick and G. Leugering, Gas flow in fan-shaped networks: Classical solutions and feedback stabilization, *SIAM J. Control Optim.*, **49** (2011), 2101–2117.
- [10] M. Gugat and M. Sigalotti, Stars of vibrating strings: Switching boundary feedback stabilization, *Netw. Heterog. Media*, **5** (2010), 299–314.
- [11] V. T. Haimo, Finite time controllers, *SIAM J. Control Optim.*, **24** (1986), 760–770.
- [12] V. Komornik, *Exact Controllability and Stabilization. The Multiplier Method*, RAM: Research in Applied Mathematics, Masson, Paris; John Wiley & Sons, Ltd., Chichester, 1994.
- [13] J. E. Lagnese, G. Leugering and E. J. P. G. Schmidt, *Modeling, analysis and control of dynamic elastic multi-link structures*, Systems & Control: Foundations & Applications, Birkhäuser Boston, Inc., Boston, MA, 1994.
- [14] A. Majda, Disappearing solutions for the dissipative wave equation, *Indiana Univ. Math. J.*, **24** (1974/75), 1119–1133.

- [15] E. Moulay and W. Perruquetti, [Finite-time stability and stabilization: state of the art](#), in *Advances in Variable Structure and Sliding Mode Control*, Lecture Notes in Control and Inform. Sci., 334, Springer, Berlin, 2006, 23–41.
- [16] V. Perrollaz and L. Rosier, Finite-time stabilization of hyperbolic systems over a bounded interval, in *1st IFAC workshop on Control of Systems Governed by Partial Differential Equations (CPDE2013)*, 2013, 239–244.
- [17] V. Perrollaz and L. Rosier, [Finite-time stabilization of  \$2 \times 2\$  hyperbolic systems on tree-shaped networks](#), *SIAM J. Control Optim.*, **52** (2014), 143–163.
- [18] Y. Shang, D. Liu and G. Xu, [Super-stability and the spectrum of one-dimensional wave equations on general feedback controlled networks](#), *IMA J. Math. Control and Inform.*, **31** (2014), 73–99.
- [19] J. Valein and E. Zuazua, [Stabilization of the wave equation on 1-D networks](#), *SIAM J. Control Optim.*, **48** (2009), 2771–2797.
- [20] Y. Zhang and G. Xu, [Controller design for bush-type 1-D wave networks](#), *ESAIM Control Optim. Calc. Var.*, **18** (2012), 208–228.
- [21] Y. Zhang and G. Xu, [Exponential and super stability of a wave network](#), *Acta Appl. Math.*, **124** (2013), 19–41.
- [22] G. Q. Xu, D. Y. Liu and Y. Q. Liu, [Abstract second order hyperbolic system and applications to controlled network of strings](#), *SIAM J. Control Optim.*, **47** (2008), 1762–1784.

Received October 2014; revised January 2015.

*E-mail address:* [fatiha.alabau@univ-lorraine.fr](mailto:fatiha.alabau@univ-lorraine.fr)

*E-mail address:* [Vincent.Perrollaz@lmpt.univ-tours.fr](mailto:Vincent.Perrollaz@lmpt.univ-tours.fr)

*E-mail address:* [Lionel.Rosier@mines-paristech.fr](mailto:Lionel.Rosier@mines-paristech.fr)